



800-656-7625

fax 610-367-7825

Product Specification Sheet

Product: IRDye™ 700DX Conjugated Affinity Purified anti-Myc EPITOPE TAG (Rabbit)

Code: 600-430-381

Lot # 14887

Size: 100 µg

Physical State: Lyophilized

Antibody Concentration: 1.0 mg/ml (by UV absorbance at 280 nm)

Label: IRDye™ 700DX (MW 1954)

Fluorochrome/Protein Ratio: 3.0 IRDye™ 700DX per mole of Rabbit IgG

Absorption Wavelength: 689 nm

Emission Wavelength: 700 nm

Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Stabilizer: 10 mg/ml BSA IgG and Protease free

Preservative: 0.01% (w/v) Sodium Azide

Background Information: Epitope tags are short peptide sequences that are easily recognized by tag-specific antibodies. Due to their small size, epitope tags do not affect the tagged protein's biochemical properties. Most often sequences encoding the epitope tag are included with target DNA at the time of cloning to produce fusion proteins containing the epitope tag sequence. This allows anti-epitope tag antibodies to serve as universal detection reagents for any tag containing protein produced by recombinant means. This means that anti-epitope tag antibodies are a useful alternative to generating specific antibodies to identify, immunoprecipitate or immunoaffinity purify a recombinant protein. The anti-epitope tag antibody is usually functional in a variety of antibody-dependent experimental procedures. Expression vectors producing epitope tag fusion proteins are available for a variety of host expression systems including bacteria, yeast, insect and mammalian cells. Rockland Immunochemicals produces anti-epitope tag antibodies against many common epitope tags including Myc, GST, GFP, 6X His, MBP, FLAG and HA. Rockland Immunochemicals also produces antibodies to other tags including FITC, Rhodamine (TRITC), DNP and biotin.

Application(s): Fluorescence technology is widely used to detect proteins. However, many common visible fluorophores often result in considerable background fluorescence in the visible range. Visible fluorophores are rarely used for membrane-based protein detection because of this high background. IRDye™ 800 and IRDye™ 700DX antibody and reagent conjugates are specifically designed for protein detection methods that use longer-wavelength, near-infrared (IR) fluorophores to visualize proteins in western blotting and other applications. Very low background fluorescence in the IR range provides for a much higher signal-to-noise ratio than visible fluorophores. Detection levels in the picogram range on Western blots rival the sensitivity of chemiluminescence on film. IRDye™ 800 conjugates are optimized for the Odyssey® Infrared Imaging System developed by LI-COR. IRDye™ 800 conjugates are also suitable for immunofluorescence microscopy using commercially available excitation/emission filters in the 780nm/820nm range. Dual simultaneous labeling in western blots or microscopy is achieved when IRDye™ 800 conjugates are used in conjunction with IRDye™ 700 or Cy5.5™ conjugates. IRDye™ 800 and IRDye™ 700DX conjugates provide an ultra-sensitive and convenient alternative to standard chemiluminescent protein detection methods, as well as a valuable tool for multicolor imaging. Anti-myc has utility to detect the fusion protein of the myc epitope cloned along with the target gene. As such, anti-myc/myc can be used to identify fusion proteins containing the myc epitope. The antibody recognizes the Myc tag fused either to the amino- or carboxy- termini of targeted proteins.

Recommended Dilution(s): This product was tested by immunoblot using Myc tagged recombinant protein spotted to nitrocellulose membrane. A 1:2,500 dilution is sufficient to detect 12-25 pg of immobilized protein. Researchers should determine optimal titers for other applications.

Storage Conditions: Store vial at 4° C prior to restoration. Restore with 0.1 ml of deionized water (or equivalent). This product is stable for several weeks at 4° C as an undiluted liquid. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. Dilute only prior to immediate use. Expiration date is one (1) year from date of restoration.

Purity and Specificity: This affinity-purified antibody is directed against myc epitope tag and is useful in determining its presence in various assays. This polyclonal anti-Myc-tag antibody detects over-expressed proteins containing the Myc epitope tag. The antibody recognizes the Myc-tag (Glu-Gln-Lys-Leu-Ile-Ser-Glu-Glu-Asp-Leu) fused to either the amino- or carboxy-termini of targeted proteins in transfected or transformed cells.

Immunogen: This antibody was purified from whole rabbit serum prepared by repeated immunizations with Myc epitope tag peptide E-Q-K-L-I-S-E-E-D-L conjugated to KLH using maleimide.

Conjugation Reference: LI-COR Biosciences, Lincoln, NE.

Related Product(s):

#600-401-381	Affinity Purified Anti-Myc TAG (Rabbit)
#600-401-382	Affinity Purified Anti-6X His TAG (Rabbit)
#600-401-383	Affinity Purified Anti-FLAG (RABBIT)
#600-401-384	Affinity Purified Anti-HEMAGGLUTININ (RABBIT)
#200-101-150	IgG Fraction of Anti-LUCIFERASE (Firefly) GOAT
#600-101-200	Affinity Purified Anti-GST (GOAT)
#600-101-215	Affinity Purified Anti-GFP (GOAT)
#200-4136	IgG Fraction of Anti-BETA GALACTOSIDASE (E.coli) (RABBIT)
#600-101-098	Affinity Purified Anti-BIOTIN (GOAT)
#600-101-096	Affinity Purified Anti-FITC (GOAT)
#200-301-246	Protein A Purified Mouse Mab Anti-RHODAMINE
#200-B01-380	Protein A Purified Hamster Mab Anti-DNP
#611-703-127	HRP Anti-Rabbit IgG [H&L] MX10 (DONKEY)
#611-132-122	IRDye800 Anti-Rabbit IgG [H&L] MX10 (GOAT)

USDA Certification: All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation.

Note: This material is subject to proprietary rights and is sold under license from LI-COR, Inc. This product is licensed for sale only for 'research-use' only. There is no implied license hereunder for any commercial use. IRDye is a trademark of LI-COR, Inc. COMMERCIAL USE shall include:

1. Resale, lease, license or other transfer of the material or any material derived or produced from it.
2. Resale, lease, license or other grant of rights to use this material or any material derived or produced from it.
3. Use of this material to perform services for a fee for third parties.

If you require a commercial license to use this material and do not have one, return this material, unopened to Rockland Inc. PO BOX 326, Gilbertsville, PA and money paid for the material will be refunded.