

## Product Specification Sheet

**Product:** Biotin Conjugated IgG fraction of anti-Asparaginase [*E. coli*] [Rabbit]

**Code:** 200-4671

**Lot #** 5410

**Size:** 10.0 mg

**Physical State:** Lyophilized

**Antibody Concentration:** 10.0 mg/ml (by UV absorbance at 280 nm)

**Label:** Biotinamidocaproate N-Hydroxysuccinimide Ester (BAC)

**Biotin/Protein Ratio:** 10-20 BAC molecules per Rabbit IgG molecule

**Buffer:** 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

**Stabilizer:** 10 mg/ml Bovine Serum Albumin (BSA) IgG and Protease free

**Preservative:** 0.01% (w/v) Sodium Azide

**Application(s):** immunoblotting, ELISA, immunohistochemistry, immunomicroscopy as well as other antibody based assays using streptavidin or avidin conjugates requiring lot-to-lot consistency.

**Recommended Dilution(s):** This product has been assayed against 1.0 ug of Asparaginase in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:3,200 to 1:12,800 of the reconstitution concentration is suggested for this product. Optimal titers for other applications should be determined by the researcher.

**Storage Conditions:** Store vial at 4° C prior to restoration. Restore with 1.0 ml of deionized water (or equivalent). For extended storage reconstitute product with 50% glycerol instead of water and then aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. Expiration date is one (1) year from date of restoration.

**Purity and Specificity:** This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin, anti-Rabbit Serum as well as purified and partially purified Asparaginase [*E. coli*]. Cross reactivity against Asparaginase from other sources may occur but have not been specifically determined.

**Immunogen:** Asparaginase [*E. coli*]

**Conjugation Reference:** Bayer & Wilchek *Methods in Enzymology* **184**; 138-160, 1990.

**Note:** This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information.