

Product Specification Sheet

Product: Affinity Purified Anti-Laminin (Human) [Rabbit] Minimum Cross Reactivity to collagens and non-collagen extracellular matrix proteins.

Code: 600-401-116-0.1

Lot # 9708

Size: 0.1 mg

Physical State: Liquid (sterile filtered)

Antibody Concentration: 1.0 mg/ml (by UV absorbance at 280 nm)

Buffer: 0.125M Sodium Borate, 0.075M Sodium Chloride, 0.005M EDTA; pH 8.0

Stabilizer: None

Preservative: 0.01% (w/v) Sodium Azide

Application(s): Anti-Laminin antibodies have been used for indirect trapping ELISA for quantitation of antigen in serum using a standard curve, for immunoprecipitation and for western blotting for highly sensitive qualitative analysis.

Recommended Dilution(s): This product was assayed by immunoblot and found to be reactive against Laminin at a dilution of 1:5,000 to 1:10,000. This product was also assayed against 1.0 µg of Laminin in a standard sandwich ELISA using Peroxidase conjugated Affinity Purified anti-Rabbit IgG [H&L] (Goat) code #611-1302 and (ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:4,000 to 1:8,000 of the stock concentration is suggested for this product. For immunohistochemistry on paraffin embedded tissue dilute the product 1:50 to 1:200. Optimal titers for other applications should be determined by the researcher.

Storage Conditions: Store vial at 4° C prior to opening. This product is stable 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage mix with an equal volume of glycerol, aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Expiration date is one (1) year from date of opening product.

Purity and Specificity: This product has been prepared by immunoaffinity chromatography using immobilized human placental laminin followed by extensive cross-adsorption against human serum proteins and collagen and non-collagen extracellular matrix proteins to remove any unwanted specificities. Typically less than 1% cross reactivity against other extracellular matrix proteins was detected by ELISA against purified standards. This antibody reacts with most mammalian Laminins and has negligible cross-reactivity with Type I, II, III, IV, V or VI Collagens or Fibronectin. Non-specific cross reaction of anti-Laminin antibodies with other human serum proteins or non-Laminin extracellular matrix proteins is negligible.

Immunogen: Produced against a mixture of synthetic peptides corresponding to regions of adherence on laminin.

Note: This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information.