

Product Specification Sheet

Product: IRDye™ 800 Conjugated Affinity Purified anti-Golden Syrian/Armenian Hamster IgG [H&L] [Goat] Minimum Cross Reactivity to Mouse and Rat Serum Proteins

Code: 620-132-440

Lot # 12833

Size: 0.5 mg

Physical State: Lyophilized

Antibody Concentration: 1.0 mg/ml (by UV absorbance at 280 nm)

Label: IRDye™ 800 (MW 1067)

Fluorochrome/Protein Ratio: 3.3 IRDye™ 800 per mole of Goat IgG

Absorption Wavelength: 778 nm

Emission Wavelength: 806 nm

Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Stabilizer: 10 mg/ml Bovine Serum Albumin (BSA) IgG and Protease free

Preservative: 0.01% (w/v) Sodium Azide

Application(s): Fluorescence technology is widely used to detect proteins. However, many common visible fluorophores often result in considerable background fluorescence in the visible range. Visible fluorophores are rarely used for membrane-based protein detection because of this high background. IRDye™ 800 antibody and reagent conjugates are specifically designed for protein detection methods that use longer-wavelength, near-infrared (IR) fluorophores to visualize proteins in western blotting and other applications. Very low background fluorescence in the IR range provides for a much higher signal-to-noise ratio than visible fluorophores. Detection levels in the picogram range rivals the sensitivity of chemiluminescence on film. IRDye™ 800 conjugates are optimized for the Odyssey® Infrared Imaging System developed by LI-COR. IRDye™ 800 conjugates are also suitable for immunofluorescence microscopy using commercially available excitation/emission filters in the 780nm/820nm range. Dual simultaneous labeling in western blots or microscopy is achieved when IRDye™ 800 conjugates are used in conjunction with Cy5.5™ conjugates. IRDye™ 800 conjugates provide an ultra-sensitive and convenient alternative to standard chemiluminescent protein detection methods, as well as a valuable tool for multicolor imaging.

Recommended Dilution(s): This product was tested by immunoblot using Armenian and Golden Syrian Hamster IgG spotted to nitrocellulose membrane. A 1:2,500 dilution is sufficient to detect 12-25 pg of immobilized IgG. Researchers should determine optimal titers for other applications.

Storage Conditions: Store vial at 4° C prior to restoration. Restore with 0.5 ml of deionized water (or equivalent). For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. Expiration date is one (1) year from date of restoration.

Purity: This product was prepared from monospecific antiserum by immunoaffinity chromatography using Hamster IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Armenian/Golden Syrian Hamster IgG and Armenian/Golden Syrian Hamster Serum. No reaction was observed against Mouse or Rat Serum Proteins.

Immunogen: Armenian/Golden Syrian Hamster IgG, whole molecule

Conjugation Reference: LI-COR Biosciences, Lincoln, NE.

Note: This material is subject to proprietary rights and is sold under license from LI-COR, Inc. This product is licensed for sale only for 'research-use' only. There is no implied license hereunder for any commercial use. IRDye is a trademark of LI-COR, Inc. COMMERCIAL USE shall include:

1. Resale, lease, license or other transfer of the material or any material derived or produced from it.
2. Resale, lease, license or other grant of rights to use this material or any material derived or produced from it.
3. Use of this material to perform services for a fee for third parties.

If you require a commercial license to use this material and do not have one, return this material, unopened to Rockland Inc. PO BOX 326, Gilbertsville, PA and money paid for the material will be refunded.